

G-1™ PLUS

Vitrolife

US

EN: Indication for use

Medium for culture of embryos from the pronucleate stage to day 2 or 3.

Product Description

SUPPLEMENTED WITH HSA
G-1™ PLUS is a bicarbonate buffered medium containing human serum albumin, hyaluronan and gentamicin as an antibacterial agent.

Ready to use after equilibration at +37 °C and 6 % CO₂ atmosphere.

Storage instructions and stability

Store dark at +2 to +8 °C.

G-1™ PLUS is stable until the expiry date shown on the container labels and the LOT-specific Certificate of Analysis.

For non-EU: Media bottles should not be stored after opening. Discard excess media after completion of the procedure.

EU: Media bottles can be used for up to two weeks after first opening. Use aseptic technique and minimize the time outside the refrigerator. Record the opening date on the bottle. Discard excess media no later than two weeks after first opening.

Directions for use

Culture of Cleavage Stage Embryos

All culture dishes (wells or tubes) should be rinsed with G-RINSE™ and prepared in advance of the procedure.

In the afternoon of the day of oocyte retrieval, label 60 mm dishes with the ID of the patient. It is recommended that dishes of the highest quality be used. Using a pre-rinsed sterilized dish, add 5 µl of G-1™ PLUS to each well. Add G-1™ PLUS into the dish. Tips are rinsed by taking up the volume required (25 µl), and then expelling this volume into a media discard dish. Four drops should be at the 3, 6, 9 and 12 o'clock positions (for embryo culture), the remaining volume is used to wash the oocytes (wash drops). Immediately cover dish with OVIDIL™ to avoid evaporation. Prepare no more than 2 dishes at one time. Using a new tip for each drop, first rinse the tip and then add a further 25 µl of medium to each original drop. This technique allows the droplets to stand up rather than flatten out.

Immediately place the dish in the incubator at 6 % CO₂ and +37 °C. Place the lid of the dish and set at an angle on the side of the plate. This will ensure proper equilibration of the dish.

Dishes must equilibrate in the incubator with a semi-opened lid for a minimum of 6 hrs (this is the minimal measured time for the media to reach correct pH under oil) and for a maximum of 18 hours.

Following removal of the cumulus cells and assessment of fertilisation, pronucleate embryos are transferred to a 1-well dish and washed in pre-warmed G-MOPS™ PLUS/ supplemented G-MOPS™ PLUS. Washing entails picking up the embryo 2-3 times in a minimal volume and moving it around the well.

Embryos should then be washed successively in the two control drops in the culture dish and up to 5 embryos placed in each 50 µl drop of G-1™ PLUS.

More than five embryos should not be cultured in each drop since it may result in a significant depletion of the nutrient pool. As a security precaution, prepare two culture dishes if the patient has more than 10 embryos. Return the dish to the incubator immediately. It is advisable to culture embryos in a group of 3 or 4. For a patient with 6 embryos it is best to culture in 2 groups of 3 rather than 4 and 2 or 5 and 1. On day 2 or on day 3, embryos can be transferred to the uterus in EmbryoGle® or in G-2™ PLUS supplemented G-2™. Alternatively, on day 3, embryos can be transferred to G-2™ PLUS/ supplemented G-2™ for further culture to the blastocyst stage.

Specifications

Sterile filtered SAL 10³

Mouse Embryo Assay (1-cell) [% expanded blastocyst within 96 hours] ≥ 80

Bacterial endotoxins (LAL assay) [EU/ml] < 0.25

LOT specific test results are available on the Certificate of Analysis provided with each delivery.

Precautions

Discard product if bottle integrity is compromised. Do not use G-1™ PLUS if it appears cloudy.

G-1™ PLUS contains human serum albumin, hyaluronan and gentamicin.

Caution: All blood products should be treated as potentially infectious. Source material from which this product was derived was found negative when tested for antibodies to HIV, HBC, HCV and HTLV III and non-reactive for HbsAg, HCV RNA and HIV-1 RNA and syphilis. No known test methods can offer assurance that products derived from human blood will not transmit infectious agents.

To avoid contamination Vitrolife strongly recommends that media should be opened and used only with aseptic technique.

The risk of reproductive toxicity and developmental toxicity for IVF media, including Vitrolife's IVF media, have not been determined and are uncertain.

Not for injection.

Caution: (EU) US law restricts this device to sale by or on the order of a physician.

Description of ISO Symbols

STERILE A Sterilized using aseptic processing techniques

10³ CFU Temperature limit

Use by - see label.

Caution: Consult accompanying documents.

REF Catalog number

LOT Batch code

CE 2460 CE Mark (Conformité Européenne)

規格

过滤器

10³

SAL

细菌繁殖(1-細胞鼠胚)

≥ 80

96小時擴張張量%

< 0.25

細菌毒素(LAL檢測)

< 0.25

每個批號的檢驗報告隨貨提供。

注意事項

EN: Indication for use

Medium for culture of embryos from the pronucleate stage to day 2 or 3.

Product Description

SUPPLEMENTED WITH HSA

G-1™ PLUS is a bicarbonate buffered medium containing human serum albumin, hyaluronan and gentamicin as an antibacterial agent.

Ready to use after equilibration at +37 °C and 6 % CO₂ atmosphere.

Storage instructions and stability

Store dark at +2 to +8 °C.

G-1™ PLUS is stable until the expiry date shown on the container labels and the LOT-specific Certificate of Analysis.

For non-EU: Media bottles should not be stored after opening. Discard excess media after completion of the procedure.

EU: Media bottles can be used for up to two weeks after first opening. Use aseptic technique and minimize the time outside the refrigerator. Record the opening date on the bottle. Discard excess media no later than two weeks after first opening.

Directions for use

Culture of Cleavage Stage Embryos

All culture dishes (wells or tubes) should be rinsed with G-RINSE™ and prepared in advance of the procedure.

In the afternoon of the day of oocyte retrieval, label 60 mm dishes with the ID of the patient. It is recommended that dishes of the highest quality be used. Using a pre-rinsed sterilized dish, add 5 µl of G-1™ PLUS to each well. Add G-1™ PLUS into the dish. Tips are rinsed by taking up the volume required (25 µl), and then expelling this volume into a media discard dish. Four drops should be at the 3, 6, 9 and 12 o'clock positions (for embryo culture), the remaining volume is used to wash the oocytes (wash drops). Immediately cover dish with OVIDIL™ to avoid evaporation. Prepare no more than 2 dishes at one time. Using a new tip for each drop, first rinse the tip and then add a further 25 µl of medium to each original drop. This technique allows the droplets to stand up rather than flatten out.

Immediately place the dish in the incubator at 6 % CO₂ and +37 °C. Place the lid of the dish and set at an angle on the side of the plate. This will ensure proper equilibration of the dish.

Dishes must equilibrate in the incubator with a semi-opened lid for a minimum of 6 hrs (this is the minimal measured time for the media to reach correct pH under oil) and for a maximum of 18 hours.

Following removal of the cumulus cells and assessment of fertilisation, pronucleate embryos are transferred to a 1-well dish and washed in pre-warmed G-MOPS™ PLUS/ supplemented G-MOPS™ PLUS. Washing entails picking up the embryo 2-3 times in a minimal volume and moving it around the well.

Embryos should then be washed successively in the two control drops in the culture dish and up to 5 embryos placed in each 50 µl drop of G-1™ PLUS.

More than five embryos should not be cultured in each drop since it may result in a significant depletion of the nutrient pool. As a security precaution, prepare two culture dishes if the patient has more than 10 embryos. Return the dish to the incubator immediately. It is advisable to culture embryos in a group of 3 or 4. For a patient with 6 embryos it is best to culture in 2 groups of 3 rather than 4 and 2 or 5 and 1. On day 2 or on day 3, embryos can be transferred to the uterus in EmbryoGle® or in G-2™ PLUS supplemented G-2™. Alternatively, on day 3, embryos can be transferred to G-2™ PLUS/ supplemented G-2™ for further culture to the blastocyst stage.

Specifications

Sterile filtered

SAL 10³

Mouse Embryo Assay (1-cell)

% expanded blastocyst within 96 hours] ≥ 80

Bacterial endotoxins (LAL assay)

[EU/ml] < 0.25

每个批号的检验证报告随货提供。

注意事項

EN: Indication for use

Medium for culture of embryos from the pronucleate stage to day 2 or 3.

Product Description

SUPPLEMENTED WITH HSA

G-1™ PLUS is a bicarbonate buffered medium containing human serum albumin, hyaluronan and gentamicin as an antibacterial agent.

Ready to use after equilibration at +37 °C and 6 % CO₂ atmosphere.

Storage instructions and stability

Store dark at +2 to +8 °C.

G-1™ PLUS is stable until the expiry date shown on the container labels and the LOT-specific Certificate of Analysis.

For non-EU: Media bottles should not be stored after opening. Discard excess media after completion of the procedure.

EU: Media bottles can be used for up to two weeks after first opening. Use aseptic technique and minimize the time outside the refrigerator. Record the opening date on the bottle. Discard excess media no later than two weeks after first opening.

Directions for use

Culture of Cleavage Stage Embryos

All culture dishes (wells or tubes) should be rinsed with G-RINSE™ and prepared in advance of the procedure.

In the afternoon of the day of oocyte retrieval, label 60 mm dishes with the ID of the patient. It is recommended that dishes of the highest quality be used. Using a pre-rinsed sterilized dish, add 5 µl of G-1™ PLUS to each well. Add G-1™ PLUS into the dish. Tips are rinsed by taking up the volume required (25 µl), and then expelling this volume into a media discard dish. Four drops should be at the 3, 6, 9 and 12 o'clock positions (for embryo culture), the remaining volume is used to wash the oocytes (wash drops). Immediately cover dish with OVIDIL™ to avoid evaporation. Prepare no more than 2 dishes at one time. Using a new tip for each drop, first rinse the tip and then add a further 25 µl of medium to each original drop. This technique allows the droplets to stand up rather than flatten out.

Immediately place the dish in the incubator at 6 % CO₂ and +37 °C. Place the lid of the dish and set at an angle on the side of the plate. This will ensure proper equilibration of the dish.

Dishes must equilibrate in the incubator with a semi-opened lid for a minimum of 6 hrs (this is the minimal measured time for the media to reach correct pH under oil) and for a maximum of 18 hours.

Following removal of the cumulus cells and assessment of fertilisation, pronucleate embryos are transferred to a 1-well dish and washed in pre-warmed G-MOPS™ PLUS/ supplemented G-MOPS™ PLUS. Washing entails picking up the embryo 2-3 times in a minimal volume and moving it around the well.

Embryos should then be washed successively in the two control drops in the culture dish and up to 5 embryos placed in each 50 µl drop of G-1™ PLUS.

More than five embryos should not be cultured in each drop since it may result in a significant depletion of the nutrient pool. As a security precaution, prepare two culture dishes if the patient has more than 10 embryos. Return the dish to the incubator immediately. It is advisable to culture embryos in a group of 3 or 4. For a patient with 6 embryos it is best to culture in 2 groups of 3 rather than 4 and 2 or 5 and 1. On day 2 or on day 3, embryos can be transferred to the uterus in EmbryoGle® or in G-2™ PLUS supplemented G-2™. Alternatively, on day 3, embryos can be transferred to G-2™ PLUS/ supplemented G-2™ for further culture to the blastocyst stage.

Specifications

Sterile filtered

SAL 10³

Mouse Embryo Assay (1-cell)

% expanded blastocyst within 96 hours] ≥ 80

Bacterial endotoxins (LAL assay)

[EU/ml] < 0.25

每个批号的检验证报告随货提供。

注意事項

EN: Indication for use

Medium for culture of embryos from the pronucleate stage to day 2 or 3.

Product Description

SUPPLEMENTED WITH HSA

G-1™ PLUS is a bicarbonate buffered medium containing human serum albumin, hyaluronan and gentamicin as an antibacterial agent.

Ready to use after equilibration at +37 °C and 6 % CO₂ atmosphere.

Storage instructions and stability

Store dark at +2 to +8 °C.

G-1™ PLUS is stable until the expiry date shown on the container labels and the LOT-specific Certificate of Analysis.

For non-EU: Media bottles should not be stored after opening. Discard excess media after completion of the procedure.

EU: Media bottles can be used for up to two weeks after first opening. Use aseptic technique and minimize the time outside the refrigerator. Record the opening date on the bottle. Discard excess media no later than two weeks after first opening.

Directions for use

Culture of Cleavage Stage Embryos

All culture dishes (wells or tubes) should be rinsed with G-RINSE™ and prepared in advance of the procedure.

In the afternoon of the day of oocyte retrieval, label 60 mm dishes with the ID of the patient. It is recommended that dishes of the highest quality be used. Using a pre-rinsed sterilized dish, add 5 µl of G-1™ PLUS to each well. Add G-1™ PLUS into the dish. Tips are rinsed by taking up the volume required (25 µl), and then expelling this volume into a media discard dish. Four drops should be at the 3, 6, 9 and 12 o'clock positions (for embryo culture), the remaining volume is used to wash the oocytes (wash drops). Immediately cover dish with OVIDIL™ to avoid evaporation. Prepare no more than 2 dishes at one time. Using a new tip for each drop, first rinse the tip and then add a further 25 µl of medium to each original drop. This technique allows the droplets to stand up rather than flatten out.

Immediately place the dish in the incubator at 6 % CO₂ and +37 °C. Place the lid of the dish and set at an angle on the side of the plate. This will ensure proper equilibration of the dish.

Dishes must equilibrate in the incubator with a semi-opened lid for a minimum of 6 hrs (this is the minimal measured time for the media to reach correct pH under oil) and for a maximum of 18 hours.

G-1™ PLUS è un terreno con tampone di bicarbonato contenente albumina del siero umano, hyaluronan e gentamicina come agente antibatterico.

Destinato all'uso dopo l'equilibratura a +37 °C in atmosfera a 6% CO₂.

Istruzioni per la conservazione e stabilità

Conservare al riparo dalla luce ad una temperatura da +2 a +8 °C.

G-1™ PLUS rimane stabile fino alla data di scadenza riportata sulle etichette dei contenitori e sul Certificato di analisi di ogni Lotto.

Non per EU: Dopo l'apertura, il terreno di terreno non possono essere conservati. Gettare il terreno in eccesso al termine della procedura.

EU: Dopo la prima apertura è possibile utilizzare i flaconi di terreno per un periodo di due settimane. Utilizzare una tecnica asettica e ridurre al minimo il tempo di permanenza dell'elemento. Seguire le dati di apertura sui flaconi. Smaltire il terreno restante al massimo due settimane dopo la prima apertura.

Istruzioni per l'uso

Coltura degli embrioni in base di clivaggio

Tutte le piastre di coltura (pocketz o provette) devono essere risciacquate con G-RINSE™ e preparate anticipatamente rispetto alla procedura.

Il pomeriggio del giorno della raccolta degli ovociti, etichettare le piastre da 60 mm con l'ID della paziente. Si ricomincia la coltura degli embrioni dopo aver rimossi una puntina sterile pre-risciacquata, posizionare silenzioso 6 x 25 µl, goccia di G-1™ supplemento/G-1™ PLUS nella piastra. Le punte sono risciacquate prelevando il volume richiesto (25 µl) e quindi espellendo tale volume in una piastra di eliminazione di due settimane. Quattro gocce di coltura si aggiungono alla piastra di coltura (piastre). È possibile avviare le piastre di coltura (gocce di lavaggio). Coprire immediatamente le gocce con OVOIL™ per evitare l'evaporazione. Non preparare più di 2 piastre per volta. Utilizzando una nuova puntina per ogni goccia, risciacquare prima la puntina poi aggiungere altri 25 µl di terreno ad ogni goccia originale. Questo consente alle gocce di restare verticali anziché appiattite.

Posizionare immediatamente la piastra nell'incubatore a 6% CO₂ e a +37 °C. Rimuovere delicatamente il copricapi della piastra e mettere in posizione angolata sul lato della piastra. Ciò garantisce la corretta equilibratura della piastra.

Le piastre devono equilibrarsi nell'incubatore con copricapi semipermeabile per un minimo di 6 ore (questo è il tempo minimo misurato perché i terreni raggiungano la maturità).

Dopo l'equilibratura, le piastre sono destinate alla fecondazione, trasferire gli embrioni prenucleati in una piastra mono-pozzetto e lavorarli in G-MOPS™ PLUS pre-risciacquo o G-MOPS™ supplemento. Per il lavaggio, prelevare 2-3 volte gli embrioni in un minimo volume e muoverli all'interno del pozzetto.

Lavare successivamente gli embrioni nelle due gocce centrali della piastra di coltura e posizionare fino a 5 embrioni in ciascuna delle gocce da 50 µl di G-1™ PLUS.

In ogni goccia può essere messa a coltura un massimo di cinque embrioni, assicurando di puoi verificare una riduzione significativa del pool di embrioni.

Ciò consente di preparare due piastre di coltura da se la paziente ha più di 10 embrioni. Ripetere immediatamente la piastra nell'incubatore. Si raccomanda di mettere in coltura gli embrioni alimento in gruppi di 2. Ad esempio, per una paziente con 6 embrioni è preferibile mettere in coltura 3 embrioni per la prima parte di 2 e 2 per la seconda parte di 1. Il giorno 2, gli embrioni possono essere trasferiti nell'utero in EmbryoGlo® oppure in G-2™ PLUS / G-2™ supplemento. In alternativa, il giorno 3, gli embrioni possono essere trasferiti in G-2™ PLUS / G-2™ supplemento per la successiva coltura alla fase blastocisti.

Specifiche

Filtraggio sterile

SAL 10³

Analisi su embrione di topo (1 cellula)

(% di blastocisti espansa entro 96 ore)

≥ 80

Endotossine batteriche (analisi LAL)

[EU/ml]

< 0.25

I risultati dei test specifici sui Lotti sono riportati sul Certificato di analisi fornito ad ogni consegna.

Precauzioni

Scartare il prodotto se l'integrità della bottiglia è compromessa. Non utilizzare G-1™ PLUS se presenta un aspetto torbido.

G-1™ PLUS contiene albumina del siero umano, hyaluronan e gentamicina.

Attenzione: gli embrioni devono essere trattati come materie prime infettive. La matrice prima impiegata per questi prodotti sono risultate negative ai test per gli anticorpi di HIV, Hbc, Hcv e H1v RNA e sifilide. Nessun metodo di prova consoluto può assicurare che i prodotti derivati dal sangue umano non trasmettano agenti infettivi.

Per evitare contaminazioni, Vitriforte raccomanda di utilizzare il prodotto esclusivamente con tecniche asettiche.

I rischi di tossicità riproduttiva e tossicità dello sviluppo del terreno (IVF), inclusi i terreni IVF Vitriforte, non sono stati determinati e sono incerti.

Non per iniezione.

Attenzione: la legge federale (degli Stati Uniti) limita la vendita del presente dispositivo dietro prescrizione medica.

LT: Naudojimo indikacija

Terpe embrionais auginti nuogios stadijos 1k-2osios ar 3-tosios dienos.

Produktu aprašas

PAPIŁDYZTA HSA

G-1™ PLUS yra bikarbonato buferinė terpė su žmogaus serumu albuminiu, hyaluronan ir gentamicinu kaip antikaterinke veikliajai medžiagai.

Parengta naudoti su pusiausvyrus nusistovėjimo esant +37 °C temperatūrą ir 6% CO₂ aplinkoje.

Laikymo nurodymai ir stabilumas

Laikyti temperatuose nuo +2 iki +40 °C.

G-1™ PLUS yra stabili iki galiojamų datos, nurodytos ant taiplykių etiketėje ar senijos analizės.

Nuo 6 iki 12 mėnesių.

Atspalvės skaidrumas 0.15-0.25.

Antspalvės skaidrumas 0.15-0.25.

<